

# THERMAL DECOMPOSITION OF CYCLOHEXANE BY FLASH PYROLYSIS VACUUM ULTRAVIOLET PHOTOIONIZATION TIME-OF-FLIGHT MASS SPECTROMETRY: A STUDY ON THE INITIAL UNIMOLECULAR DECOMPOSITION MECHANISM

KUANLIANG SHAO, *Department of Chemistry, University of California, Riverside, Riverside, CA, USA*; XINGHUA LIU, *School of Science, Hainan University, Haikou, Hainan, China*; PAUL JONES, GE SUN, *Department of Chemistry, University of California, Riverside, Riverside, CA, USA*; MARIAH GOMEZ, *Department of Chemistry and Biochemistry, University of California, Los Angeles, Los Angeles, CA, USA*; BLAKE RISER, *Department of Biochemistry, University of California, Riverside, Riverside, CA, USA*; JING-SONG ZHANG, *Department of Chemistry, University of California, Riverside, Riverside, CA, USA*.

Thermal decomposition of cyclohexane at temperatures up to 1310 K was performed using flash pyrolysis coupled with vacuum ultraviolet (118.2 nm) photoionization time-of-flight mass spectrometry. The experimental results revealed that the major initiation reaction of cyclohexane decomposition was C-C bond fission leading to the formation of 1,6-hexyl diradical. The 1,6-hexyl diradical could isomerize to 1-hexene and decompose into  $\bullet\text{C}_3\text{H}_7 + \bullet\text{C}_3\text{H}_5$  and  $\bullet\text{C}_4\text{H}_7 + \bullet\text{C}_2\text{H}_5$ . The 1,6-hexyl diradical could also undergo direct dissociation; the  $\text{C}_4\text{H}_8$  fragment via the 1,4-butyl diradical intermediate was observed, serving as evidence of the 1,6-hexyl diradical mechanism. Quantum chemistry calculations at UCCSD(T)/cc-pVDZ level of theory on the initial reaction pathways of cyclohexane were performed and found to be consistent with the experimental conclusions. Cyclohexyl radical was not observed as an initial intermediate in the pyrolysis. Benzene was produced from sequential  $\text{H}_2$  eliminations of cyclohexane at high temperatures.